

ASPECTS OF DYNAMICAL CHIRAL SYMMETRY BREAKING

C. D. ROBERTS

*Physics Division, Argonne National Laboratory
Argonne, Illinois, 60439-4843, USA*

Dynamical chiral symmetry breaking is a nonperturbative phenomenon that may be studied using QCD's gap equation. Model-independent results can be obtained with a nonperturbative and symmetry preserving truncation. The gap equation yields the massive dressed-quark propagator, which has a spectral representation when considered as a function of the current-quark mass. This enables an explanation of the connection between the infrared limit of the QCD Dirac operator's spectrum and the quark condensate appearing in the operator product expansion.

1. Introduction

Dynamical chiral symmetry breaking (DCSB) is fundamental to the strong interaction spectrum. It is the generation through self-interactions of a momentum-dependent quark mass, $M(p^2)$, in the chiral limit, that is large in the infrared but power-suppressed in the ultraviolet:

$$M(p^2) \stackrel{\text{large-}p^2}{=} \frac{2\pi^2\gamma_m}{3} \frac{(-\langle\bar{q}q\rangle^0)}{p^2 \left(\frac{1}{2} \ln \left[\frac{p^2}{\Lambda_{\text{QCD}}^2}\right]\right)^{1-\gamma_m}}. \quad (1)$$

The phenomenon is impossible in weakly interacting theories. In Eq. (1), $\gamma_m = 12/(33 - 2N_f)$ is the mass anomalous dimension, with N_f the number of light-quark flavours, and $\langle\bar{q}q\rangle^0$ is the renormalisation-group-invariant vacuum quark condensate.¹ Dynamical chiral symmetry breaking is an essentially nonperturbative effect.

It is natural to explore DCSB using QCD's gap equation and the evolution of the dressed-quark mass-function in Eq. (1) to a large and finite constituent-quark-like mass in the infrared, $M(0) \sim 0.5 \text{ GeV}$, is a long-standing prediction of such Dyson-Schwinger equation (DSE) studies² that has recently been confirmed in simulations of quenched lattice-QCD.³ The features and flaws of the approach are well known. For example, the DSEs are a keystone in proving renormalisability and provide a generating tool for

perturbation theory. However, they are also a collection of coupled integral equations from which a tractable problem is only obtained by truncation. A weak coupling expansion, which reproduces perturbation theory, is one systematic truncation procedure. However, that cannot be used to study nonperturbative phenomena.

2. Truncating the DSEs

Fortunately a systematic alternative is available.⁴ Its leading order term furnishes a reliable description of vector and flavour nonsinglet pseudoscalar mesons and their interactions,^{5,6} and the reasons for success in these channels can be understood *a priori*. The scheme is worth illustrating.

The renormalised gap equation

$$S(p)^{-1} = Z_2 (i\gamma \cdot p + m^{\text{bm}}) + Z_1 \int_q^\Lambda g^2 D_{\mu\nu}(p-q) \frac{\lambda^a}{2} \gamma_\mu S(q) \Gamma_\nu^a(q, p), \quad (2)$$

involves: $D_{\mu\nu}(k)$, the dressed-gluon propagator; $\Gamma_\nu^a(q; p)$, the dressed-quark-gluon vertex; m^{bm} , the Λ -dependent current-quark bare mass that appears in the Lagrangian; $\int_q^\Lambda := \int^\Lambda d^4q / (2\pi)^4$, a translationally-invariant regularisation of the integral, with Λ the regularisation mass-scale; and the quark-gluon-vertex and quark wave function renormalisation constants, $Z_1(\zeta^2, \Lambda^2)$ and $Z_2(\zeta^2, \Lambda^2)$ respectively, which depend on the renormalisation point, the regularisation mass-scale and the gauge parameter.

The kernel of Eq. (2) is formed from a product of the dressed-gluon propagator and dressed-quark-gluon vertex. It is to this product that a systematic truncation must be applied. However, one must also consider more than just the gap equation's kernel. Chiral symmetry is expressed via the axial-vector Ward-Takahashi identity:

$$P_\mu \Gamma_{5\mu}(k; P) = S^{-1}(k_+) i\gamma_5 + i\gamma_5 S^{-1}(k_-), \quad (3)$$

$k_\pm = k \pm P/2$, wherein $\Gamma_{5\mu}(k; P)$ is the dressed axial-vector vertex. This three-point function satisfies an inhomogeneous Bethe-Salpeter equation:

$$[\Gamma_{5\mu}(k; P)]_{tu} = Z_2 [\gamma_5 \gamma_\mu]_{tu} + \int_q^\Lambda [S(q_+) \Gamma_{5\mu}(q; P) S(q_-)]_{sr} K_{tu}^{rs}(q, k; P), \quad (4)$$

in which $K(q, k; P)$ is the renormalised fully-amputated quark-antiquark scattering kernel, and the colour-, Dirac- and flavour-matrix structure in the equation is denoted by the indices r, s, t, u . The Ward-Takahashi identity, Eq. (3), means that the kernel in the gap equation and that in the Bethe-Salpeter equation are intimately related. Therefore a qualitatively reliable

understanding of chiral symmetry and its dynamical breaking can only be obtained using a truncation scheme that preserves this relation, and hence guarantees Eq. (3) without a *fine-tuning* of model-dependent parameters.

The truncation scheme introduced in Ref. [4] is a dressed-loop expansion of the dressed-quark-gluon vertices that appear in the half-amputated dressed-quark-antiquark scattering matrix: S^2K , a renormalisation-group invariant.⁵ All n -point functions involved thereafter in connecting two particular quark-gluon vertices are *fully dressed*. The effect of this truncation in the gap equation, Eq. (2), is realised through the following representation of the quark-gluon vertex, $\Gamma_\mu = \frac{1}{2}\lambda^a \Gamma_\mu$:

$$\begin{aligned} Z_1 \Gamma_\mu(k, p) = & \gamma_\mu + \frac{1}{2N_c} \int_\ell^\Lambda g^2 D_{\rho\sigma}(p - \ell) \gamma_\rho S(\ell + k - p) \gamma_\mu S(\ell) \gamma_\sigma \\ & + \frac{N_c}{2} \int_\ell^\Lambda g^2 D_{\sigma'\sigma}(\ell) D_{\tau'\tau}(\ell + k - p) \gamma_{\tau'} S(p - \ell) \gamma_{\sigma'} \Gamma_{\sigma\tau\mu}^{3g}(\ell, -k, k - p) \\ & + [\dots]. \end{aligned} \quad (5)$$

Here Γ^{3g} is the dressed-three-gluon vertex and it is apparent that the lowest order contribution to each term written explicitly is $O(g^2)$. The ellipsis represents terms whose leading contribution is $O(g^4)$; viz., the crossed-box and two-rung dressed-gluon ladder diagrams, and also terms of higher leading-order.

This expansion of S^2K , with its implications for other n -point functions, yields an ordered truncation of the DSEs that guarantees, term-by-term, the preservation of vector and axial-vector Ward-Takahashi identities, a feature that has been exploited^{7,8} to prove Goldstone's theorem and other exact results in QCD. It is readily seen that inserting Eq. (5) into Eq. (2) provides the rule by which the rainbow-ladder truncation can be systematically improved.

With this scheme, as with perturbation theory, it is impossible, in general, to obtain complete closed-form expressions for the kernels of the gap and Bethe-Salpeter equations. However, for the planar dressed-quark-gluon vertex depicted in Fig. 1, closed forms can be obtained and a number of significant features illustrated⁵ when one uses the following model for the dressed-gluon line⁹

$$g^2 D_{\mu\nu}(k) = \left[\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right] (2\pi)^4 G^2 \delta^4(k), \quad (6)$$

where G sets the model's mass-scale. This model has many positive features and, furthermore, its particular momentum-dependence works to advantage

$$\Gamma_\mu^a(k, p) = \text{[Diagram 1]} + \text{[Diagram 2]} + \text{[Diagram 3]} + \dots$$

Figure 1. Integral equation for a planar dressed-quark-gluon vertex obtained by neglecting contributions associated with explicit gluon self-interactions. Solid circles indicate fully dressed propagators. The vertices are not dressed. (Adapted from Ref. [5].)

in reducing integral equations to character-preserving algebraic equations. Naturally, there is a drawback: the simple momentum dependence also leads to some model-dependent artefacts, but they are easily identified and hence not cause for concern.

It is a general result⁵ that with any vertex whose diagrammatic content is known explicitly; e.g., Fig. 1, it is possible to construct a unique Bethe-Salpeter kernel which ensures the Ward-Takahashi identities are fulfilled: that kernel is necessarily nonplanar. This becomes transparent with the model in Eq. (6), using which the gap equation obtained with the vertex depicted in Fig. 1 reduces to an algebraic equation, irrespective of the number of dressed-gluon rungs that are retained, and the same is true of the Bethe-Salpeter equations in every channel: pseudoscalar, vector, etc.

Results for the π and ρ are illustrated in Table 1. It is apparent that, irrespective of the order of the truncation; i.e., the value of n , the number of dressed gluon rungs in the quark-gluon vertex, the pion is massless in the chiral limit. (NB. This pion is composed of heavy dressed-quarks, as is evident in the calculated scale of the dynamically generated dressed-quark mass function: $M(0) \approx 0.5 \text{ GeV}$.) The masslessness of the π is a model-independent consequence of consistency between the Bethe-Salpeter kernel and the kernel in the gap equation. Furthermore, the bulk of the

Table 1. Calculated π and ρ masses, in GeV, quoted with $G = 0.48 \text{ GeV}$. n is the number of dressed-gluon rungs retained in the planar vertex, see Fig. 1, and hence the order of the vertex-consistent Bethe-Salpeter kernel: the rapid convergence of the kernel is evident from the tabulated results. (Adapted from Ref. [5].)

	$M_H^{n=0}$	$M_H^{n=1}$	$M_H^{n=2}$	$M_H^{n=\infty}$
$\pi, m = 0$	0	0	0	0
$\pi, m = 0.011$	0.152	0.152	0.152	0.152
$\rho, m = 0$	0.678	0.745	0.754	0.754
$\rho, m = 0.011$	0.695	0.762	0.770	0.770

ρ - π mass splitting is present in the chiral limit and with the simplest ($n = 0$; i.e., rainbow-ladder) kernel, which shows that this mass difference is driven by the DCSB mechanism: it is not the result of a finely adjusted hyperfine interaction. Finally, the quantitative effect of improving on the rainbow-ladder truncation; i.e., including more dressed-gluon rungs in the gap equation's kernel and consistently improving the kernel in the Bethe-Salpeter equation, is a 10% correction to the vector meson mass. Simply including the first correction ($n = 1$; i.e., retaining the first two diagrams in Fig. 1) gives a vector meson mass that differs from the fully resummed result by $\lesssim 1\%$.

While these results were obtained with a rudimentary interaction model, the procedure is completely general. However, the algebraic simplicity of the analysis is naturally peculiar to the model. With a more realistic interaction, the gap and vertex equations yield a system of twelve coupled integral equations. The Bethe-Salpeter kernel for any given channel then follows as the solution of a determined integral equation.

By identifying the rainbow-ladder truncation as the lowest order in a systematic scheme the procedure also provides a means of anticipating the channels in which that truncation must fail. The scalar mesons are an example. Parametrisations of the rainbow-ladder truncation, fitted to π , ρ observables, yield scalar mesons masses that are too large.¹⁰ That was thought to be a problem. However, we now know this had to happen because cancellations that occur between higher order terms in the pseudoscalar and vector channels, thereby reducing the magnitude of corrections, do not occur in the scalar channel, wherein the full kernel contains additional attraction.¹¹ Therefore an interaction model employed in rainbow-ladder truncation which simultaneously provides a good description of scalar and pseudoscalar mesons must contain spurious degrees of freedom. Quantitative studies of the effect of the higher-order terms have begun^{11,12} and a straightforward understanding of scalar mesons is not impossible. Similarly, it may be that the only thing remarkable about mesons with “exotic” quantum numbers is that their accurate description requires only an equally carefully considered application of this systematic approach.¹³

3. Concerning the quark condensate

While Eq. (1) is expressed in Landau gauge, the so-called OPE condensate, $\langle \bar{q}q \rangle^0$, is gauge parameter independent. In the chiral limit this condensate sets the scale of the mass function in the ultraviolet and thus plays

a role analogous to that of the renormalisation-group-invariant current-quark mass in the massive theory. A determination of the OPE condensate directly from lattice-QCD simulations must await an accurate chiral extrapolation¹⁴ but DSE models tuned to reproduce modern lattice data give¹⁵ $|\langle \bar{q}q \rangle^0| \sim \Lambda_{\text{QCD}}^3$.

Another view of DCSB is obtained by considering the eigenvalues and eigenfunctions of the anti-Hermitian massless Euclidean Dirac operator:

$$\gamma \cdot D u_n(x) = i\lambda_n u_n(x). \quad (7)$$

The eigenfunctions form a complete set, and except for zero modes they occur in pairs: $\{u_n(x), \gamma_5 u_n(x)\}$, with eigenvalues of opposite sign. It follows that in an external gauge field, A , one can write the quark Green function

$$S(x, y; A) = \langle q(x) \bar{q}(y) \rangle_A^m = \sum_n \frac{u_n(x) u_n^\dagger(y)}{i\lambda_n + m}, \quad (8)$$

where m is the current-quark mass. Assuming, e.g., a lattice regularisation,

$$\frac{1}{V} \int_V d^4x \langle \bar{q}(x) q(x) \rangle_A^m = -\frac{2m}{V} \sum_{\lambda_n > 0} \frac{1}{\lambda_n^2 + m^2}, \quad (9)$$

where V is the lattice volume.^a One may now define a quark condensate:

$$\langle 0 | \bar{q}q | 0 \rangle^m := \lim_{V \rightarrow \infty} \frac{1}{V} \int_V d^4x \langle \bar{q}(x) q(x) \rangle_A^m, \quad (10)$$

wherein the r.h.s. expresses an average over gauge field configurations. When $V \rightarrow \infty$ the operator spectrum becomes dense and Eqs. (9), (10) become

$$-\langle 0 | \bar{q}q | 0 \rangle^m = 2m \int_0^\infty d\lambda \frac{\rho(\lambda)}{\lambda^2 + m^2}, \quad (11)$$

with $\rho(\lambda)$ a spectral density. This equation expresses an assumption that in QCD the full two-point massive-quark Schwinger function, when viewed as a function of the current-quark mass, has a spectral representation.

It follows formally from Eq. (11) that

$$-\langle 0 | \bar{q}q | 0 \rangle^0 := \lim_{m \rightarrow 0} 2m \int_0^\infty d\lambda \frac{\rho(\lambda)}{\lambda^2 + m^2} = \pi \rho(0). \quad (12)$$

This is the so-called Banks-Casher relation,¹⁷ which has long been advocated as a means by which a quark condensate may be measured in lattice-QCD simulations.¹⁸

^aIn deriving Eq. (9), zero modes have been neglected, which is justified under broad conditions.¹⁶

The explication¹⁹ of a correspondence between the condensate in Eq. (1) and that in Eq. (12) requires care because Eq. (11) is meaningless as written: dimensional counting reveals the r.h.s. has mass-dimension three and since λ will at some point be greater than any relevant internal scale, the integral must diverge as Λ^2 , where Λ is the regularising mass-scale.

The Schwinger function

$$\tilde{\sigma}(m) := N_c \text{tr}_D \int_p^\Lambda \tilde{S}_m(p), \quad (13)$$

which is the trace of the unrenormalised massive dressed-quark propagator evaluated at a fixed value of the regularisation scale, Λ , can be identified with the l.h.s. of Eq. (11). The essence of the Banks-Casher relation is that $\tilde{\sigma}(m)$ have a spectral representation:

$$\tilde{\sigma}(m) := 2m \int_0^\Lambda d\lambda \frac{\tilde{\rho}(\lambda)}{\lambda^2 + m^2}, \quad (14)$$

where $m = m^{\text{bm}}(\Lambda)$, and Eq. (14) entails

$$\tilde{\rho}(\lambda) = \frac{1}{2\pi} \lim_{\eta \rightarrow 0^+} [\tilde{\sigma}(i\lambda + \eta) - \tilde{\sigma}(i\lambda - \eta)]. \quad (15)$$

The content of this sequence of equations is elucidated by inserting the free quark propagator in Eq. (13). The integral thus obtained is readily evaluated using dimensional regularisation:

$$\tilde{\sigma}_{\text{free}}(m) = \frac{N_c}{4\pi^2} m^3 \left[\ln \frac{m^2}{\zeta^2} + \frac{1}{\varepsilon} + \gamma - \ln 4\pi \right]. \quad (16)$$

With Eq. (15) the regularisation dependent terms cancel and one obtains

$$\tilde{\rho}(\lambda) = \frac{N_c}{4\pi^2} \lambda^3. \quad (17)$$

In perturbation theory it can be shown that every contribution to $\tilde{\rho}(\lambda)$ is proportional to λ^3 and hence

$$\langle 0 | \bar{q}q | 0 \rangle^0 \propto \tilde{\rho}(\lambda = 0) = 0. \quad (18)$$

A nonzero value of $\rho(0)$ is an essentially nonperturbative effect.

A precise analysis requires that attention be paid to renormalisation. Therefore consider the gauge-parameter-independent trace of the renormalised massive quark propagator:

$$\sigma(m; \zeta) = Z_4(\zeta, \Lambda) N_c \text{tr}_D \int_p^\Lambda S_m(p; \zeta),$$

where the argument remains $m = m^{\text{bm}}(\Lambda)$. The renormalisation constant Z_4 vanishes logarithmically with increasing Λ and hence one still has $\sigma(m) \sim \Lambda^2 m^{\text{bm}}(\Lambda)$. However, since⁷

$$-\langle \bar{q}q \rangle_\zeta^0 = \lim_{\Lambda \rightarrow \infty} Z_4(\zeta, \Lambda) N_c \text{tr}_D \int_p^\Lambda S_0(p; \zeta), \quad (19)$$

it is clear that for any finite but large value of Λ and tolerance δ , it is always possible to find $m_\delta^{\text{bm}}(\Lambda)$ such that

$$\sigma(m; \zeta) + \langle \bar{q}q \rangle_\zeta^0 < \delta, \quad \forall m^{\text{bm}} < m_\delta^{\text{bm}}. \quad (20)$$

This is true in QCD and can be illustrated using the model of Ref. [1]. Figure 2 displays $\sigma(m, \zeta)$, evaluated using a hard cutoff, Λ , on the integral in Eq. (19), calculated with the massive dressed-quark propagators obtained by solving the gap equation. Since Eq. (20) specifies the domain on which the value of $\sigma(m; \zeta)$ is determined by nonperturbative effects, one anticipates

$$m_\delta^{\text{bm}}(\Lambda) \approx -\langle \bar{q}q \rangle_\zeta^0 / \Lambda^2 \sim 10^{-9} \quad (21)$$

for $\Lambda = 2.0 \text{ TeV}$ in QCD where $|\langle \bar{q}q \rangle_\zeta^0| \sim \Lambda_{\text{QCD}}^3$. This estimate is confirmed in Fig. 2.

The dotted line in Fig. 2 is

$$\sigma(m, \zeta) = -\langle \bar{q}q \rangle_\zeta^0 \frac{2}{\pi} \arctan \frac{\Lambda}{m} + Z_4(\zeta, \Lambda) \frac{N_c}{4\pi^2} m [\Lambda^2 - m^2 \ln(1 + \Lambda^2/m^2)], \quad (22)$$

and the difference between this and the curve is of order $\alpha(\Lambda) m(\Lambda) \Lambda^2$ because the DSE model incorporates QCD's one-loop behaviour. The figure also displays $\sigma(m, \zeta)$ obtained in the absence of confinement, in which case²⁰ $\langle \bar{q}q \rangle_\zeta^0 \equiv 0$, as is apparent.

It is now plain that in QCD $\sigma(m, \zeta)$ is a monotonically increasing convex-up function with a regular chiral limit and consequently it has a spectral representation:

$$\sigma(m, \zeta) = 2m \int_0^\Lambda \frac{\rho(\lambda)}{\lambda^2 + m^2}. \quad (23)$$

This lays the vital plank in a veracious connection between the condensates in Eqs. (1) and (12). On the domain specified by Eq. (21), the behaviour of $\sigma(m, \zeta)$ in Eq. (19) is given by Eq. (22), which yields, via Eq. (15),

$$\pi \rho(\lambda) = -\langle \bar{q}q \rangle_\zeta^0 + Z_4(\zeta, \Lambda) \frac{N_c}{4\pi} \lambda^3 + \dots, \quad (24)$$




Figure 2. Circles/solid-line: $\sigma(m)^{1/3}$ in Eq. (19) as a function of the current-quark bare-mass, evaluated using the dressed-quark propagator obtained in the model of Ref. [1]; dashed line: the model's value of $(-\langle \bar{q}q \rangle_{\zeta=1 \text{ GeV}}^0) = (0.24 \text{ GeV})^3$; and dotted line: Eq. (22). Diamonds: $\sigma(m)^{1/3}$ evaluated in a non-confining version of the model; dot-dashed line: Eq. (22) with $\langle \bar{q}q \rangle^0 \equiv 0$. (Adapted from Ref. [19].)

where the ellipsis denotes contributions from the higher-order terms implicit in Eq. (22).

Reference [21] reports the spectral density of the staggered Dirac operator in quenched $SU(3)$ gauge theory calculated on a $V = 4^4$ -lattice, in the vicinity of the deconfining phase transition at $\beta \gtrsim 5.6$. A comparison¹⁹ with these results shows that while the effect of finite lattice volume is evident early (for $\lambda a \gtrsim 0.1$, a is the lattice spacing) the behaviour at small λa is qualitatively in agreement with Eq. (24) and Fig. 2: a nonzero OPE condensate dominates the Dirac spectrum in the confined domain; and it vanishes in the deconfined domain, whereupon $\rho(0) = 0$ and the perturbative evolution, Eq. (17), is manifest. To be more quantitative, for a lattice coupling $\beta = 5.4$, $\rho(0)a^3 \approx 70$, so that

$$\pi \rho(0) a^3 / V \approx (0.95)^3. \quad (25)$$

The lattice spacing was not determined in Ref. [21] but one can nevertheless assess the scale of Eq. (25) by supposing $a \sim 0.3 \text{ fm} \sim 0.3/\Lambda_{\text{QCD}}$, a value not unreasonable for small lattices and small β , wherewith the r.h.s. is

$\sim (3\Lambda_{\text{QCD}})^3$. This is too large but not unreasonable given the parameters of the simulation, its errors and the systematic uncertainties in our estimate. Fitting the lattice data at $\beta = 5.8$, one finds $\rho(\lambda) \propto \lambda^3$ on $\lambda < 0.1$ but with a proportionality constant larger than that anticipated from perturbation theory; viz. Eq. (24). Some mismatch was to be expected, however, because at $\beta = 5.8$ one has only just entered the deconfined domain and near the transition boundary some nonperturbative effects are still relevant. It is a modern challenge to determine those gauge couplings and lattice parameters for which the data are quantitatively consistent with Eq. (24).

In practice, there are three main parameters in a simulation of lattice-QCD: the lattice volume, characterised by a length L ; the lattice spacing, a ; and the current-quark mass, m . So long as the lattice size is large compared with the current-quark's Compton wavelength; viz., $L \gg 1/m$, then dynamical chiral symmetry breaking can be expressed in the simulation. Supposing that to be the case then, as outlined above, so long as the lattice spacing is small compared with the current-quark's Compton wavelength

$$\pi \rho(0) \approx -\langle \bar{q}q \rangle_{1/a}^0, \quad a \ll 1/m \ll L, \quad (26)$$

where the $\langle \bar{q}q \rangle_{1/a}^0$ is the scale-dependent OPE condensate [$\zeta = \Lambda = 1/a$ in Eq. (24)]. This completes an explication of the connection between the condensates. Note, however, that the continuum analysis indicates that one requires $am \lesssim (a\Lambda_{\text{QCD}})^3$ if $\rho(\lambda = 0)$ is to provide a veracious estimate of the OPE condensate. The residue at the lowest-mass pole in the flavour-nonsinglet pseudoscalar vacuum polarisation provides a measure of the OPE condensate that is accurate for larger current-quark masses.¹⁹

Acknowledgments

I am grateful to the Organising Committee for the opportunity to participate in the “5th International Conference on Quark Confinement and the Hadron Spectrum.” This work was supported by: the Department of Energy, Nuclear Physics Division, contract no. W-31-109-ENG-38; the National Science Foundation, grant no. INT-0129236; and benefited from the resources of the National Energy Research Scientific Computing Center.

References

1. P. Maris and C.D. Roberts, *Phys. Rev.* **C56**, 3369 (1997).
2. C.D. Roberts and A.G. Williams, *Prog. Part. Nucl. Phys.* **33**, 477 (1994).
3. P.O. Bowman, U.M. Heller and A.G. Williams, *Phys. Rev.* **D66**, 014505 (2002).

4. A. Bender, C.D. Roberts and L. v. Smekal, *Phys. Lett.* **B380**, 7 (1996).
5. A. Bender, W. Detmold, A.W. Thomas and C.D. Roberts, *Phys. Rev.* **C65**, 065203 (2002).
6. P. Maris and C.D. Roberts, “Dyson-Schwinger Equations: A tool for Hadron Physics,” nucl-th/0301049.
7. P. Maris, C.D. Roberts and P.C. Tandy, *Phys. Lett.* **B420**, 267 (1998).
8. M.A. Ivanov, Yu.L. Kalinovsky and C.D. Roberts, *Phys. Rev.* **D60**, 034018 (1999).
9. H.J. Munczek and A.M. Nemirovsky, *Phys. Rev.* **D28**, 181 (1983).
10. P. Jain and H.J. Munczek, *Phys. Rev.* **D48**, 5403 (1993); P. Maris, C.D. Roberts, S.M. Schmidt and P.C. Tandy, *Phys. Rev.* **C63**, 025202 (2001); P. Maris, *Few Body Syst.* **32**, 41 (2002).
11. S.R. Cotanch and P. Maris, “QCD based quark description of $\pi - \pi$ scattering up to the σ and ρ region,” hep-ph/0210151.
12. A. Höll, P. Maris and C.D. Roberts, *Phys. Rev.* **C59**, 1751 (1999); J.C.R. Bloch, C.D. Roberts and S.M. Schmidt, *Phys. Rev.* **C60**, 065208 (1999); P. Maris and P.C. Tandy, *Phys. Rev.* **C62**, 055204 (2000); A. Bender, W. Detmold and A.W. Thomas, *Phys. Lett.* **B516**, 54 (2001).
13. J.C.R. Bloch, Yu.L. Kalinovsky, C.D. Roberts and S.M. Schmidt, *Phys. Rev.* **D60**, 111502 (1999); C.J. Burden and M.A. Pichowsky, *Few Body Syst.* **32**, 119 (2002).
14. J.B. Zhang, F.D.R. Bonnet, P.O. Bowman, D.B. Leinweber and A.G. Williams, “Towards the Continuum Limit of the Overlap Quark Propagator in Landau Gauge,” hep-lat/0208037.
15. P. Maris, A. Raya, C.D. Roberts and S.M. Schmidt, “Confinement and dynamical chiral symmetry breaking,” nucl-th/0208071.
16. H. Leutwyler and A. Smilga, *Phys. Rev.* **D46**, 5607 (1992).
17. T. Banks and A. Casher, *Nucl. Phys.* **B169**, 103 (1980).
18. E. Marinari, G. Parisi and C. Rebbi, *Phys. Rev. Lett.* **47**, 1795 (1981).
19. K. Langfeld, R. Pullirsch, H. Markum, C.D. Roberts and S.M. Schmidt, “Concerning the quark condensate,” nucl-th/0301024.
20. A. Bender, D. Blaschke, Yu.L. Kalinovsky and C.D. Roberts, *Phys. Rev. Lett.* **77**, 3724 (1996); F.T. Hawes, P. Maris and C.D. Roberts, *Phys. Lett.* **B440**, 353 (1998).
21. E. Bittner, H. Markum and R. Pullirsch, *Nucl. Phys. Proc. Suppl.* **96**, 189 (2001).